

BCS-BEC crossover with Rashba and Dresselhaus couplings

Luca Salasnich

Dipartimento di Fisica e Astronomia "Galileo Galilei", Università di Padova, Italy

Trieste, May 15, 2013


Collaboration with:
Luca Dell'Anna and Giovanni Mazarella (Università di Padova)

Summary

- BCS-BEC crossover
- Artificial spin-orbit coupling
- Mean-field approach
- Singlet and triplet condensate
- Results with Rashba coupling
- Including Dresselhaus coupling
- Conclusions
- Acknowledgments

BCS-BEC crossover (I)

In 2004 the BCS-BEC crossover has been observed with **ultracold gases made of fermionic ^{40}K and ^6Li alkali-metal atoms.**¹



This crossover is obtained by changing (with a **Feshbach resonance**) the s-wave scattering length a_s of the inter-atomic potential:

- $a_s \rightarrow 0^-$ (BCS regime of weakly-interacting Cooper pairs)
- $a_s \rightarrow \pm\infty$ (unitarity limit of strongly-interacting Cooper pairs)
- $a_s \rightarrow 0^+$ (BEC regime of bosonic dimers)

¹C.A. Regal et al., PRL **92**, 040403 (2004); M.W. Zwierlein et al., PRL **92**, 120403 (2004); M. Bartenstein, A. Altmeyer et al., PRL **92**, 120401 (2004); J. Kinast et al., PRL **92**, 150402 (2004).

BCS-BEC crossover (II)

The crossover from a BCS superfluid ($a_s < 0$) to a BEC of molecular pairs ($a_s > 0$) has been investigated experimentally around a **Feshbach resonance**, where the s-wave scattering length a_s diverges, and it has been shown that the system is (meta)stable.

The detection of **quantized vortices** under rotation² has clarified that **this dilute and ultracold gas of Fermi atoms is superfluid**.

Usually the BCS-BEC crossover is analyzed in terms of

$$y = \frac{1}{k_F a_s} \quad (1)$$

the inverse **scaled interaction strength**, where $k_F = (3\pi^2 n)^{1/3}$ is the Fermi wave number and n the total density.

The system is dilute because $r_e k_F \ll 1$, with r_e the effective range of the inter-atomic potential.

²M.W. Zwierlein *et al.*, Science **311**, 492 (2006); M.W. Zwierlein *et al.*, Nature **442**, 54 (2006).

Artificial spin-orbit coupling

In 2011 and 2012 **artificial spin-orbit coupling** has been experimentally imposed on both bosonic³ and fermionic⁴ atomic gases.

The **single-particle Hamiltonian** \hat{h}_{sp} with both **Rashba and Dresselhaus spin-orbit couplings** reads

$$\hat{h}_{sp} = \frac{\hat{p}^2}{2m} + v_R (\hat{\sigma}_x \hat{p}_y - \hat{\sigma}_y \hat{p}_x) + v_D (\hat{\sigma}_x \hat{p}_y + \hat{\sigma}_y \hat{p}_x) , \quad (2)$$

with $\hat{p}^2 = -\hbar^2 \nabla^2$, $\hat{p}_x = -i\hbar \frac{\partial}{\partial x}$, $\hat{p}_y = -i\hbar \frac{\partial}{\partial y}$, v_R and v_D the Rashba and Dresselhaus coupling constant, respectively, and

$$\hat{\sigma}_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} , \quad \hat{\sigma}_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} .$$

³Y.J. Lin, K. Jimenez-Garcia, and I.B. Spielman, Nature **471**, 83 (2011).

⁴P. Wang et al., PRL **109**, 095301 (2012); L.W. Cheuk et al., PRL **109**, 095302 (2012).

Mean-field approach (I)

The **partition function** \mathcal{Z} of the uniform two-spin-component Fermi system at temperature T , in a volume V , and with chemical potential μ can be written in terms of a functional integral as

$$\mathcal{Z} = \int \mathcal{D}[\psi_s, \bar{\psi}_s] \exp \left\{ -\frac{1}{\hbar} S \right\}, \quad (3)$$

where

$$S = \int_0^{\hbar\beta} d\tau \int_V d^3\mathbf{r} \mathcal{L} \quad (4)$$

is the **Euclidean action functional** and \mathcal{L} is the **Euclidean Lagrangian density**, given by

$$\mathcal{L} = (\bar{\psi}_\uparrow, \bar{\psi}_\downarrow) \left[\hbar\partial_\tau + \hat{h}_{sp} - \mu \right] \begin{pmatrix} \psi_\uparrow \\ \psi_\downarrow \end{pmatrix} + g \bar{\psi}_\uparrow \bar{\psi}_\downarrow \psi_\downarrow \psi_\uparrow \quad (5)$$

with g is the strength of the s-wave coupling ($g < 0$ in the BCS regime). Notice that $\beta = 1/(k_B T)$ with k_B the Boltzmann constant. In the rest of the seminar we shall use units such that $\hbar = m = k_B = 1$.

Mean-field approach (II)

The **Lagrangian density** \mathcal{L} is quartic in the fermionic fields ψ_s , but one can reduce the problem to a quadratic Lagrangian density by introducing an auxiliary complex scalar field $\Delta(\mathbf{r}, \tau)$ via **Hubbard-Stratonovich transformation**⁵, which gives

$$\mathcal{Z} = \int \mathcal{D}[\psi_s, \bar{\psi}_s] \mathcal{D}[\Delta, \bar{\Delta}] \exp \{-S_e\}, \quad (6)$$

where

$$S_e = \int_0^{1/T} d\tau \int_V d^3\mathbf{r} \mathcal{L}_e \quad (7)$$

and the (exact) **effective Euclidean Lagrangian density** \mathcal{L}_e reads

$$\mathcal{L}_e = (\bar{\psi}_\uparrow, \bar{\psi}_\downarrow) \left[\partial_\tau + \hat{h}_{sp} - \mu \right] \begin{pmatrix} \psi_\uparrow \\ \psi_\downarrow \end{pmatrix} + \bar{\Delta} \psi_\downarrow \psi_\uparrow + \Delta \bar{\psi}_\uparrow \bar{\psi}_\downarrow - \frac{|\Delta|^2}{g}. \quad (8)$$

⁵H.T.C. Stoof, K.B. Gubbels, D.B.M. Dickerscheid, *Ultracold Quantum Fields* (Springer, Dordrecht, 2009).

Mean-field approach (III)

It is a standard procedure to integrate out the quadratic fermionic fields and to get a new formally-exact **effective action** S_{eff} which depends only on the auxiliary field $\Delta(\mathbf{r}, \tau)$. In this way we obtain

$$\mathcal{Z} = \int \mathcal{D}[\Delta, \bar{\Delta}] \exp \{-S_{eff}\}, \quad (9)$$

where

$$S_{eff} = -Tr[\ln(G^{-1})] - \int_0^{1/T} d\tau \int_V d^3\mathbf{r} \frac{|\Delta|^2}{g} \quad (10)$$

with $\gamma(\hat{\mathbf{p}}) = v_R(\hat{p}_y + i\hat{p}_x) + v_D(\hat{p}_y - i\hat{p}_x)$ and

$$G^{-1} = \begin{pmatrix} \partial_\tau + \frac{\hat{p}^2}{2} - \mu & \Delta & \gamma(\hat{\mathbf{p}}) & 0 \\ \bar{\Delta} & \partial_\tau - \frac{\hat{p}^2}{2} + \mu & 0 & -\gamma(-\hat{\mathbf{p}}) \\ \bar{\gamma}(\hat{\mathbf{p}}) & 0 & \partial_\tau + \frac{\hat{p}^2}{2} - \mu & -\Delta \\ 0 & -\bar{\gamma}(-\hat{\mathbf{p}}) & -\bar{\Delta} & \partial_\tau - \frac{\hat{p}^2}{2} + \mu \end{pmatrix} \quad (11)$$

Mean-field approach (IV)

For a uniform Fermi superfluid within the **simplest mean-field approximation** one has a **constant and real gap parameter**, i.e. $\Delta(\mathbf{r}, \tau) = \Delta$, and the **partition function** becomes⁶

$$\mathcal{Z}_{mf} = \exp \{-S_{mf}\} = \exp \left\{ -\frac{\Omega_{mf}}{T} \right\}, \quad (12)$$

where

$$S_{mf} = \frac{\Omega_{mf}}{T} = - \sum_{\mathbf{k}} \left[\sum_{j=1}^4 \ln \left(1 + e^{-E_{\mathbf{k},j}/T} \right) - \frac{\xi_{\mathbf{k}}}{T} \right] - \frac{V}{T} \frac{\Delta^2}{g} \quad (13)$$

with $\xi_{\mathbf{k}} = k^2/2 - \mu$, $\gamma_{\mathbf{k}} = v_R(k_y + ik_x) + v_D(k_y - ik_x)$, and

$$E_{\mathbf{k},1} = \sqrt{(\xi_{\mathbf{k}} - |\gamma_{\mathbf{k}}|)^2 + \Delta^2}, \quad E_{\mathbf{k},3} = -E_{\mathbf{k},1}, \quad (14)$$

$$E_{\mathbf{k},2} = \sqrt{(\xi_{\mathbf{k}} + |\gamma_{\mathbf{k}}|)^2 + \Delta^2}, \quad E_{\mathbf{k},4} = -E_{\mathbf{k},2}. \quad (15)$$

⁶L. Dell'Anna, G. Mazarella, L.S., PRA **84**, 033633 (2011).

Mean-field approach (V)

The constant and real gap parameter Δ is obtained from

$$\frac{\partial S_{mf}}{\partial \Delta} = 0, \quad (16)$$

which gives the **gap equation**

$$-\frac{1}{g} = \frac{1}{V} \sum_{\mathbf{k}} \sum_{j=1,2} \frac{\tanh(E_{\mathbf{k},j}/2T)}{4E_{\mathbf{k},j}}. \quad (17)$$

The integral on the right side of this equation is formally divergent. However, expressing the bare interaction strength g in terms of the physical scattering length a_s with the formula⁷

$$-\frac{1}{g} = -\frac{1}{4\pi a_s} + \frac{1}{V} \sum_{\mathbf{k}} \frac{1}{k^2} \quad (18)$$

one obtains the **regularized gap equation**⁸

$$-\frac{1}{4\pi a_s} = \frac{1}{V} \sum_{\mathbf{k}} \left[\sum_{j=1,2} \frac{\tanh(E_{\mathbf{k},j}/2T)}{4E_{\mathbf{k},j}} - \frac{1}{k^2} \right]. \quad (19)$$

⁷M. Marini, F. Pistoiesi, G.C. Strinati, EPJ B **1**, 151 (1998).

⁸L. Dell'Anna, G. Mazarella, L.S., PRA **84**, 033633 (2011).

Mean-field approach (VI)

From the thermodynamic formula

$$N = - \left(\frac{\partial \Omega_{mf}}{\partial \mu} \right)_{V, T} \quad (20)$$

one obtains also the **equation for the number of particles**⁹

$$N = \sum_{\mathbf{k}} \left(1 - \frac{\xi_{\mathbf{k}} - |\gamma_{\mathbf{k}}|}{2E_{\mathbf{k},1}} \tanh(E_{\mathbf{k},1}/2T) - \frac{\xi_{\mathbf{k}} + |\gamma_{\mathbf{k}}|}{2E_{\mathbf{k},2}} \tanh(E_{\mathbf{k},2}/2T) \right) . \quad (21)$$

⁹L. Dell'Anna, G. Mazzarella, L.S., PRA **84**, 033633 (2011).

Singlet and triplet condensation (I)

In a Fermi system the largest eigenvalue N_C of the two-body density matrix gives the **number of correlated fermion pairs which have their center of mass with zero linear momentum**.¹⁰ This **condensed number of pairs** is given by

$$N_C = N_0 + N_1, \quad (22)$$

where

$$N_0 = \int d^3\mathbf{r} d^3\mathbf{r}' \left[|\langle \psi_{\downarrow}(\mathbf{r}) \psi_{\uparrow}(\mathbf{r}') \rangle|^2 + |\langle \psi_{\uparrow}(\mathbf{r}) \psi_{\downarrow}(\mathbf{r}') \rangle|^2 \right] \quad (23)$$

is the **condensed number of pairs in the spin 0 state** ($m_s = 0$), while

$$N_1 = \int d^3\mathbf{r} d^3\mathbf{r}' \left[|\langle \psi_{\uparrow}(\mathbf{r}) \psi_{\uparrow}(\mathbf{r}') \rangle|^2 + |\langle \psi_{\downarrow}(\mathbf{r}) \psi_{\downarrow}(\mathbf{r}') \rangle|^2 \right]. \quad (24)$$

is the **condensed number of pairs in the spin 1 state** ($|m_s| = 1$).

¹⁰A.J. Leggett, Quantum liquids. Bose condensation and Cooper pairing in condensed-matter systems (Oxford Univ. Press, Oxford, 2006).

Singlet and triplet condensation (II)

In our superfluid Fermi system with spin-orbit coupling we obtain¹¹

$$N_0 = \frac{\Delta^2}{4} \sum_{\mathbf{k}} \left(\frac{1}{2E_{\mathbf{k},1}} \tanh(E_{\mathbf{k},1}/2T) + \frac{1}{2E_{\mathbf{k},2}} \tanh(E_{\mathbf{k},2}/2T) \right)^2. \quad (25)$$

and

$$N_1 = \frac{\Delta^2}{4} \sum_{\mathbf{k}} \left(\frac{1}{2E_{\mathbf{k},1}} \tanh(E_{\mathbf{k},1}/2T) - \frac{1}{2E_{\mathbf{k},2}} \tanh(E_{\mathbf{k},2}/2T) \right)^2. \quad (26)$$

Notice that **in the absence of spin-orbit coupling** ($v_R = v_D = 0$) one has $E_{\mathbf{k},1} = E_{\mathbf{k},2}$ from which one gets $N_1 = 0$, and consequently **the condensate number of Cooper pairs in the triplet state is zero.**

¹¹L. Dell'Anna, G. Mazarella, L.S., PRA **84**, 033633 (2011).

Results with Rashba coupling (I)

We are interested in the low temperature regime where the **condensate fraction** can be quite large. Quantitatively we restrict our study to the **zero temperature limit ($T=0$)**. In the equations above we have therefore simply $\tanh(E_{\mathbf{k},j}/2T) \rightarrow 1$.

In this way the **regularized gap equation** is given by

$$-\frac{1}{4\pi a_s} = \frac{1}{V} \sum_{\mathbf{k}} \left[\sum_{j=1,2} \frac{1}{4E_{\mathbf{k},j}} - \frac{1}{k^2} \right], \quad (27)$$

while the **number equation** reads

$$N = \sum_{\mathbf{k}} \left(1 - \frac{\xi_{\mathbf{k}} - |\gamma_{\mathbf{k}}|}{2E_{\mathbf{k},1}} - \frac{\xi_{\mathbf{k}} + |\gamma_{\mathbf{k}}|}{2E_{\mathbf{k},2}} \right). \quad (28)$$

Results with Rashba coupling (II)

Similarly, we obtain for the **spin 0 condensate number**

$$N_0 = \frac{\Delta^2}{4} \sum_{\mathbf{k}} \left(\frac{1}{2E_{\mathbf{k},1}} + \frac{1}{2E_{\mathbf{k},2}} \right)^2. \quad (29)$$


and for the **spin 1 condensate number**

$$N_1 = \frac{\Delta^2}{4} \sum_{\mathbf{k}} \left(\frac{1}{2E_{\mathbf{k},1}} - \frac{1}{2E_{\mathbf{k},2}} \right)^2. \quad (30)$$

From the previous equations one can calculate the **chemical potential** μ , the **energy gap** Δ , and also the **condensate fractions** $N_0/(N/2)$ and $N_1/(N/2)$, as a function of the **scaled interaction strength** $y = 1/(k_F a_S)$.


Note: We now show the results obtained for $v_D = 0$, i.e. when **only Rashba spin-orbit coupling is active**.

Results with Rashba coupling (III)




Scaled chemical potential μ/ϵ_F as a function of the dimensional interaction strength $\gamma = 1/(k_F a_s)$ for different values of the scaled Rashba velocity: $v_R/v_F = 0$ (solid line), $v_R/v_F = 0.7$ (long-dashed line), $v_R/v_F = 1$ (short-dashed line), $v_R/v_F = 1.4$ (dotted line), $v_R/v_F = 2$ (dashed-dotted line). Here $\epsilon_F = v_F^2/2$ is the Fermi energy and $v_F = (3\pi^2 n)^{1/3}$ is the Fermi velocity.

Results with Rashba coupling (IV)



Scaled energy gap Δ/ϵ_F as a function of the dimensional interaction strength $\gamma = 1/(k_F a_s)$ for different values of the scaled Rashba velocity: $v_R/v_F = 0$ (solid line), $v_R/v_F = 0.7$ (long-dashed line), $v_R/v_F = 1$ (short-dashed line), $v_R/v_F = 1.4$ (dotted line), $v_R/v_F = 2$ (dashed-dotted line). Here $\epsilon_F = v_F^2/2$ is the Fermi energy and $v_F = (3\pi^2 n)^{1/3}$ is the Fermi velocity.

Results with Rashba coupling (V)



Spin 0 condensate fraction $n_0/(n/2)$ (upper curves) and spin 1 condensate fraction $n_1/(n/2)$ (lower curves) as a function of the adimensional interaction strength $y = 1/(k_F a_s)$ for different values of the scaled Rashba velocity: $v_R/v_F = 0$ (solid line), $v_R/v_F = 0.7$ (long-dashed line), $v_R/v_F = 1$ (short-dashed line), $v_R/v_F = 1.4$ (dotted line), $v_R/v_F = 2$ (dashed-dotted line). Here $v_F = (3\pi^2 n)^{1/3}$ is the Fermi velocity.

Including Dresselhaus coupling (I)

We now consider also the Dresselhaus coupling, i.e. $v_D \neq 0$.
For simplicity we set¹²


$$v_R = v \cos(\theta) , \quad (31)$$

$$v_D = v \sin(\theta) , \quad (32)$$

where θ is the mixing angle.


¹²L. Dell'Anna, G. Mazarella, L.S., PRA **86**, 053632 (2012).

Including Dresselhaus coupling (II)



Spin 0 condensate fraction $n_0/(n/2)$ as a function of the **dimensional interaction strength** $y = 1/(k_F a_s)$ for three values of ν and two values of θ : $\theta = 0$ means $v_R = \nu$ and $v_D = 0$ (**red solid curves**), while $\theta = \pi/4$ means $v_R = v_D = \nu/\sqrt{2}$ (**blue dotted curves**). ν is in units of the Fermi velocity $v_F = (3\pi^2 n)^{1/3}$.

Including Dresselhaus coupling (III)



Spin 1 condensate fraction $n_1/(n/2)$ as a function of the adimensional interaction strength $y = 1/(k_F a_s)$ for three values of ν and two values of θ : $\theta = 0$ means $v_R = \nu$ and $v_D = 0$ (red solid curves), while $\theta = \pi/4$ means $v_R = v_D = \nu/\sqrt{2}$ (blue dotted curves). ν is in units of the Fermi velocity $v_F = (3\pi^2 n)^{1/3}$.

Conclusions

- With a large spin-orbit coupling v the chemical potential μ is negative for any interaction strength y .
- In the deep BCS regime ($y \ll -1$) and small v we get the analytical result

$$\mu = \epsilon_F - v^2 .$$

- A finite condensate fraction n_1/n of triplet pairs appears due to the spin-orbit coupling, in addition to the familiar condensate fraction n_0/n of singlet pairs.
- In the case where Rashba is equal to Dresselhaus, i.e. $v_R = v_D = v/\sqrt{2}$ with $\theta = \pi/4$, we find the following formulas (for any y)

$$\mu(v, \theta = \pi/4) = \mu(v = 0) - v^2 ,$$

$$\Delta(v, \theta = \pi/4) = \Delta(v = 0) .$$

$$n_c(v, \theta = \pi/4) = n_c(v = 0) ,$$

where $n_c = n_0 + n_1$ is the total condensate density.

THANK YOU FOR YOUR ATTENTION!

Our results on this topic are published in

L. Dell'Anna, G. Mazzearella, and L.S., PRA **84**, 033633 (2011).

L. Dell'Anna, G. Mazzearella, and L.S., PRA **86**, 053632 (2012).

We acknowledge **research grants** from:

- Università di Padova: Progetto di Ricerca di Ateneo 2012-2013.
- Fondazione CARIPARO: Progetto di Eccellenza 2012-2013.
- MIUR: Progetto PRIN call 2011-2012.